

LOCAL STRUCTURE OF MODULI SPACES

3. Mathematische Arbeitstagung (Neue Serie)

20.-27 Juni 1997

Vadim Schechtman

Max-Planck-Institut für Mathematik,
Gottfried-Claren-Straße 26, 53225 Bonn

This is a report on a joint work with Vladimir Hinich.

1. Let X be a smooth algebraic variety over a field k of characteristic 0. Assume that X has no infinitesimal automorphisms, i.e. $H^0(X; \mathcal{T}_X) = 0$ ($\mathcal{T}_X$ being the sheaf of vector fields). Let $\mathcal{M} = \text{Spec}(\hat{\mathcal{O}}_{\mathcal{M}})$ be the formal moduli space of deformations of X ; $\hat{\mathcal{O}}_{\mathcal{M}}$ is a complete local k -algebra.

We have the *Kodaira-Spencer isomorphism*

$$T_{\mathcal{M};X} = H^1(X; \mathcal{T}_X) \quad (KS)$$

Here $T_{\mathcal{M};X} = (\mathfrak{m}_{\mathcal{O}_{\mathcal{M}}}/\mathfrak{m}_{\mathcal{O}_{\mathcal{M}}}^2)^*$ is the tangent space of $\mathcal{M}$ at X . We want to consider the

Problem. *Describe the whole algebra $\hat{\mathcal{O}}_{\mathcal{M}}$ in terms of X .*

Let

$$\mathcal{T}_X^\bullet = R\Gamma(X; \mathcal{T}_X) : 0 \longrightarrow \mathcal{T}^0 \longrightarrow \mathcal{T}^1 \longrightarrow \dots$$

be a complex computing the sheaf cohomology of $\mathcal{T}_X$. The last sheaf is a sheaf of Lie algebras, hence $\mathcal{T}^\bullet$ may be chosen to be a *differential graded Lie algebra*; it is a correctly defined object of the appropriate derived category of *Homotopy Lie Algebras*, [HS1].

Theorem 1. *One has a canonical isomorphism of k -algebras*

$$\hat{\mathcal{O}}_{\mathcal{M}} = [H_0^{Lie}(\mathcal{T}_X^\bullet)]^* \quad (1)$$

The homology of a (dg) Lie algebra is a (dg) coalgebra. The dual space is an algebra. The isomorphism (1) is a generalization of the Kodaira-Spencer isomorphism.

This theorem is a just an example of a quite general fact; the similar results (with the same proof) hold true for other deformation problems. For example, we may wish to describe deformations of group representations, etc. Cf. [S2], [HS1].

I know two proofs of Theorem 1. The first one works in the case when $\mathcal{M}$ is smooth, and uses the *higher Kodaira-Spencer maps*, cf. [HS1]. The second one works in general situation. It uses certain very natural *sheaf property of Lie-Deligne functor*, and is described below.

2. Deligne groupoids. Let $\mathfrak{g} = \oplus_{i \geq 0} \mathfrak{g}^i$ be a nilpotent dg Lie algebra. Recall that *groupoid* is a category with all morphisms being isomorphisms. The *Deligne groupoid* $\mathcal{G}(\mathfrak{g}^\bullet)$ is defined as follows. Its objects are *Maurer-Cartan elements*

$$MC(\mathfrak{g}^\bullet) := \{y \in \mathfrak{g}^1 \mid dy + \frac{1}{2}[y, y] = 0\}$$

Let $\mathcal{G}(\mathfrak{g}^0)$ be the Lie group corresponding to the nilpotent Lie algebra $\mathfrak{g}^0$. The algebra $\mathfrak{g}^0$ acts on $MC(\mathfrak{g}^\bullet)$ by the rule

$$x \circ y = dx + [x, y], \quad x \in \mathfrak{g}^0, y \in MC(\mathfrak{g}^\bullet),$$

hence the group $\mathcal{G}(\mathfrak{g}^0)$ acts on $MC(\mathfrak{g}^\bullet)$. By definition,

$$Hom_{\mathcal{G}(\mathfrak{g}^\bullet)}(y, y') = \{g \in \mathcal{G}(\mathfrak{g}^0) \mid y' = gy\}$$

Morphisms are composed in the obvious way. Of course, this is a generalization of the Lie functor from Lie algebras to Lie groups.

3. Let us return to our deformation situation. The variety X defines a functor

$$Def_X : Art_k \longrightarrow Groupoids$$

where Art_k is the category of artinian k -algebras with residue field k . Namely, $Def_X(A)$ is the groupoid whose objects are flat deformations of X over A , and morphisms are isomorphisms identical on X .

On the other hand, if $\mathfrak{g}^\bullet$ is a dg Lie algebra over k , it defines a functor

$$\mathcal{G}_{\mathfrak{g}^\bullet} : Art_X \longrightarrow Groupoids,$$

by

$$\mathcal{G}_{\mathfrak{g}^\bullet}(A) = \mathcal{G}(\mathfrak{m}_A \otimes \mathfrak{g}^\bullet)$$

where $\mathfrak{m}_A$ is the maximal ideal of A .

Example. Assume that $X = Spec(R)$ is affine. Then one sees immediately from the definitions (Grothendieck) that one has an isomorphism of functors

$$Def_X = \mathcal{G}_{\mathfrak{g}^\bullet} \tag{2}$$

where $\mathfrak{g}^\bullet = T_X = H^0(X; \mathcal{T}_X) = Der_k(R)$ considered as a dg Lie algebra concentrated in dimension 0.

Sometimes when (2) holds, the people say that the dg Lie algebra $\mathfrak{g}^\bullet$ *governs the deformations of X* .

Theorem 2. Let X be arbitrary, and (2) holds for some $\mathfrak{g}^\bullet$. If $H^0(\mathfrak{g}^\bullet) = 0$ then

$$\hat{\mathcal{O}}_{\mathcal{M}} = (H_0^{Lie}(\mathfrak{g}^\bullet))^*$$

Proof. For an arbitrary $A \in \text{Art}_K$, we have

$$\begin{aligned} \text{Hom}_{\text{Art}_K}(\hat{\mathcal{O}}_{\mathcal{M}}, A) &= \pi_0(\text{Def}_X(A)) = \pi_0(\mathcal{G}(\mathfrak{m}_A \otimes \mathfrak{g}^\bullet)) = \\ &= \text{Hom}_{\text{alg}}((H_0^{\text{Lie}}(\mathfrak{g}^\bullet))^*, A) \triangle \end{aligned}$$

4. Now, we know the Lie algebra $\mathfrak{g}^\bullet$ for affine varieties, and we want to know it for arbitrary ones, i.e. we want to glue them.

Let $\mathfrak{g}^\bullet$ be a sheaf of nilpotent Lie algebras on a topological space X . It defines a presheaf of groupoids $\mathcal{G}(\mathfrak{g}^\bullet)$,

$$U \mapsto \mathcal{G}(\Gamma(U; \mathfrak{g}^\bullet)).$$

Theorem 3, [H]. Assume that $\mathfrak{g}^\bullet$ is a sheaf in the homotopy sense, i.e. for an open covering $\mathcal{U} = \{U_i\}$, $U = \bigcup U_i$, the natural map

$$\Gamma(U; \mathfrak{g}^\bullet) \longrightarrow \check{\mathcal{C}}(\mathcal{U}; \mathfrak{g}^\bullet)$$

is quasiisomorphism. Then $\mathcal{G}(\mathfrak{g}^\bullet)$ is a sheaf (i.e. stack).

Here $\check{\mathcal{C}}(\mathcal{U}; \mathfrak{g}^\bullet)$ is the Čech complex of the covering $\mathcal{U}$.

Theorem 1 follows immediately from Theorems 2 and 3, applied to a homotopy sheaf resolution of T_X .

5. It seems that the similar statements hold true in characteristic p , or in mixed characteristics, cf. [S2]. One should work with algebras of distributions instead of Lie algebras, and with simplicial objects instead of dg objects.

References

- [H] V. Hinich, Descent of Deligne groupoids, *Int. Math. Res. Notices* **5**(1997), 223-239.
- [HS1] V. Hinich, V. Schechtman, Deformation theory and Lie algebra homology, alg-geom/9405013, *Alg. Coll.* (1997), to appear.
- [HS2] V. Hinich, V. Schechtman, On the descent of Deligne groupoids, Preprint (October 1994).
- [S1] V. Schechtman, Localization of Deligne groupoids, Preprint (May 1994).
- [S2] V. Schechtman, Letter to B.Mazur (August 1994).